



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

294. Proposed by O. L. CALLECOT, Gettysburg, S. Dak.

Find the limit of  $\sum_{n=1}^{n=\infty} \frac{2(n^2+3n+3)}{n(n+1)(n+2)(n+3)}$ .

Solution by H. V. SPUNAR, C. E., East Pittsburg, Pa., and J. SCHEFFER, A. M., Kee Mar College, Hagerstown, Md.

$$\begin{aligned} \sum_{n=1}^{n=\infty} \frac{2(n^2+3n+3)}{n(n+1)(n+2)(n+3)} &= \sum_{n=1}^{n=\infty} \left[ \frac{1}{n} - \frac{1}{n+1} + \frac{1}{n+2} - \frac{1}{n+3} \right] \\ &= \sum \frac{1}{n} - \sum \frac{1}{n+1} + \sum \frac{1}{n+2} - \sum \frac{1}{n+3} \\ &= 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} + \frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots \\ &\quad - \frac{1}{2} - \frac{1}{3} - \frac{1}{4} - \dots - \frac{1}{n} - \frac{1}{n+1} - \frac{1}{n+2} - \frac{1}{n+3} - \dots \\ &\quad + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} + \frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots \\ &\quad - \frac{1}{4} - \dots - \frac{1}{n} - \frac{1}{n+1} - \frac{1}{n+2} - \frac{1}{n+3} - \dots \\ &= 1 + \frac{1}{3} = 1\frac{1}{3}. \end{aligned}$$

Also solved by C. E. White and G. B. M. Zerr.

In order for the solution of this problem to be rigorous, the matter of convergency must be investigated. The equality

$$\sum_{n=1}^{n=\infty} \left[ \frac{1}{n} - \frac{1}{n+1} + \frac{1}{n+2} - \frac{1}{n+3} \right] = \sum_{n=1}^{n=\infty} \frac{1}{n} - \sum_{n=1}^{n=\infty} \frac{1}{n-1} + \sum_{n=1}^{n=\infty} \frac{1}{n+2} - \sum_{n=1}^{n=\infty} \frac{1}{n+3}$$

assumed to hold in the above solution is not always true.

## GEOMETRY.


323. Proposed by CHARLES GILPIN, JR., Philadelphia, Pa.

A sphere with the radius  $R$  is divided into two segments by a plane passed through it half way between the center and circumference. The smaller segment is divided into two parts by a plane passed through it at right angles to the base and cutting it half way between its center and circumference. Required the contents of the two parts of the segment.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa., and BENJAMIN F. FINKEL, Ph. D., Drury College, Springfield, Mo.

The equation of the sphere is  $x^2 + y^2 + z^2 = R^2$ . The volume of  $C-HIK$  is

$$\begin{aligned}
V_1 &= \int dx \int dy \int dz = \int_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} dx \int_0^{\sqrt{(\frac{3}{4}R^2-x^2)}} dy \int_{\frac{1}{2}R}^{\sqrt{(R^2-x^2-y^2)}} dz \\
&= \int_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} dx \int_0^{\sqrt{(\frac{3}{4}R^2-x^2)}} [\sqrt{(R^2-x^2-y^2)} - \frac{1}{2}R] dy \\
&= \int_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} 2 \left[ \frac{1}{2}y\sqrt{(R^2-x^2-y^2)} + \frac{1}{2}(R^2-x^2)\sin^{-1} \frac{y}{\sqrt{(R^2-x^2)}} - \frac{1}{2}Ry \right]_0^{\sqrt{(\frac{3}{4}R^2-x^2)}} dx \\
&= \int_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} \left[ (R^2-x^2)\cos^{-1}\left(\frac{\frac{1}{2}R}{\sqrt{(R^2-x^2)}}\right) - \frac{1}{2}R\sqrt{(\frac{3}{4}R^2-x^2)} \right] dx \\
&= \left[ (R^2x - \frac{1}{3}x^3)\cos^{-1}\left(\frac{\frac{1}{2}R}{\sqrt{(R^2-x^2)}}\right) \right. \\
&\quad \left. - \frac{1}{24}R \int \left( \frac{9R^4 - 33R^2x^2 + 16x^4}{(R^2-x^2)\sqrt{(\frac{3}{4}R^2-x^2)}} dx \right) \right]_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} \\
&= \left[ (R^2x - \frac{1}{3}x^3)\cos^{-1} \frac{\frac{1}{2}R}{\sqrt{(R^2-x^2)}} \right. \\
&\quad \left. - \frac{1}{24}R \int \left( \frac{17R^2}{\sqrt{(\frac{3}{4}R^2-x^2)}} - \frac{16x^2}{\sqrt{(\frac{3}{4}R^2-x^2)}} - \frac{8R^4}{(R^2-x^2)\sqrt{(\frac{3}{4}R^2-x^2)}} \right) dx \right]_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} \\
&= \left[ (R^2x - \frac{1}{3}x^3)\cos^{-1} \frac{\frac{1}{2}R}{\sqrt{(R^2-x^2)}} - \frac{1}{24}R \left( 11R^2\sin^{-1}\left(\frac{x}{\frac{1}{2}\sqrt{3}R}\right) \right. \right. \\
&\quad \left. \left. - 16R^2\sin^{-1} \frac{x}{\sqrt{3(R^2-x^2)}} + 8x\sqrt{(\frac{3}{4}R^2-x^2)} \right) \right]_{\frac{1}{4}\sqrt{3}R}^{\frac{1}{2}\sqrt{3}R} \\
&= R^3 \left[ \frac{1}{72}\pi - \frac{1}{64}\sqrt{3}\cos^{-1} \frac{2}{\sqrt{13}} - \frac{2}{3}\sin^{-1} \frac{1}{\sqrt{13}} + \frac{1}{16}\sqrt{3} \right].
\end{aligned}$$



The volume of  $D-HIK$  is

$$\begin{aligned}
V_2 &= 4 \int_{\frac{1}{2}R}^R dx \int_0^{\sqrt{(R^2-x^2)}} dy \int_0^{\sqrt{(R^2-x^2-y^2)}} dz - V_1 = \frac{5}{24}\pi R^3 - V_1 \\
&= R^3 \left[ \frac{\pi}{36} + \frac{1}{64}\sqrt{3}\cos^{-1}\left(\frac{2}{\sqrt{13}}\right) + \frac{2}{3}\sin^{-1}\left(\frac{1}{\sqrt{13}}\right) - \frac{1}{16}\sqrt{3} \right].
\end{aligned}$$

Also solved with different results by H. V. Spunar and J. Scheffer.